

●代表論文

1. Wajima Z, Nakajima Y, Kim C, Kobayashi N, Kadotani H, Adachi H, Inoue T, Ogawa R: I.v. compared with brachial plexus infusion of butorphanol for postoperative analgesia. *British Journal of Anaesthesia* 74, 392-395, 1995
2. Wajima Z, Shitara T, Nakajima Y, Kim C, Kobayashi N, Kadotani H, Adachi H, Ishikawa G, Kaneko K, Inoue T, Ogawa R: Comparison of continuous brachial plexus infusion of butorphanol, mepivacaine and mepivacaine-butorphanol mixtures for postoperative analgesia. *British Journal of Anaesthesia* 75, 548-551, 1995
3. Wajima Z, Shitara T, Nakajima Y, Kim C, Kobayashi N, Kadotani H, Adachi H, Ishikawa G, Kaneko K, Inoue T, Ogawa R: Continuous brachial plexus infusion of butorphanol-mepivacaine mixtures for analgesia after upper extremity surgery. *British Journal of Anaesthesia* 78, 83-85, 1997
4. Wajima Z, Hua X-Y, Yaksh TL: Inhibition of spinal protein kinase C blocks substance P-mediated hyperalgesia. *Brain Research* 877, 314-321, 2000
5. Wajima Z, Yoshikawa T, Ogura A, Imanaga K, Shiga T, Inoue T, Ogawa R: The effects of diltiazem on hemodynamics and seizure duration during electroconvulsive therapy. *Anesthesia and Analgesia* 92, 1327-1330, 2001
6. Wajima Z, Yoshikawa T, Ogura A, Imanaga K, Shiga T, Inoue T, Ogawa R: Intravenous colforsin daropate, a water-soluble forskolin derivative, prevents thiamylal-fentanyl-induced bronchoconstriction in humans. *Critical Care Medicine* 30, 820-826, 2002
7. Wajima Z, Yoshikawa T, Ogura A, Imanaga K, Shiga T, Inoue T, Ogawa R: Oral tizanidine, an α_2 -adrenoceptor agonist, reduces minimum alveolar concentration (MAC) of sevoflurane in human adults. *Anesthesia and Analgesia* 95, 393-396, 2002
8. Wajima Z, Yoshikawa T, Ogura A, Imanaga K, Shiga T, Inoue T, Ogawa R: Intravenous verapamil blunts hyperdynamic responses during electroconvulsive therapy without altering seizure activity. *Anesthesia and Analgesia* 95, 400-402, 2002
9. Wajima Z, Yoshikawa T, Ogura A, Imanaga K, Shiga T, Inoue T, Ogawa R: Intravenous nicorandil prevents thiamylal-fentanyl-induced bronchoconstriction in humans. *Critical Care Medicine* 31, 485-490, 2003
10. Wajima Z, Shiga T, Yoshikawa T, Ogura A, Imanaga K, Inoue T, Ogawa R: Effect of prophylactic bronchodilator treatment with intravenous colforsin daropate, a water-soluble forskolin derivative, on airway resistance after tracheal intubation. *Anesthesiology* 99, 18-26, 2003
11. Wajima Z, Shiga T, Yoshikawa T, Ogura A, Imanaga K, Inoue T, Ogawa R: Intravenous alprostadil, an analog of prostaglandin E1, prevents thiamylal-fentanyl-induced bronchoconstriction in humans. *Anesthesia and Analgesia* 97, 456-460, 2003
12. Wajima Z, Shiga T, Imanaga K, Inoue T, Ogawa R: Effect of prophylactic bronchodilator treatment with i.v. carperitide on airway resistance and lung compliance after tracheal intubation. *British Journal of Anaesthesia* 96, 660-4, 2006

●著作・総説

1. 輪嶋善一郎: 緊急麻酔(小川龍編),克誠堂出版,東京,1993 担当部分:眼科手術の緊急麻酔(71-72)
2. 輪嶋善一郎: 麻酔の How to 技術編(小川龍編),克誠堂出版,東京,2001 担当部分:心機能(118-121)
3. 輪嶋善一郎: 痛みの治療薬 その基礎から臨床まで(監修 井上哲夫,鈴木勉,高橋秀則,長瀬博,成田年,益田律子,輪嶋善一郎),エルゼビア・ジャパン株式会社,東京,2005

翻訳・監修:輪嶋善一郎 第 30 章 末梢神経にオピオイドを作用させて得る鎮痛法(491-506)

監修:輪嶋善一郎 第 32 章 疼痛下における炎症性サイトカインの役割—疼痛治療への可能性(517-538)

翻訳・監修:輪嶋善一郎 第 34 章 急性痛に対する鎮痛薬の使用(563-590)

監修:輪嶋善一郎 第 35 章 鎮痛薬の相互作用(591-620)

4. 輪嶋善一郎 第 11 章 集中治療室と手術室における経頭蓋超音波ドプラ法 p102-110. 監訳 小松徹、藤原祥裕、益田律子 周術期超音波診断・治療ガイド. エルゼビア・ジャパン株式会社 東京 2010



●その他, 麻酔学会での活動, 海外派遣, 社会活動など

- ・公益社団法人日本麻酔科学会 学術集会企画専門部会
第 8 麻酔全般ワーキンググループ サテライトメンバー
- ・日本ペインクリニック学会 評議員
- ・第 4 回 Evidence-based Anesthesia 研究会 学術集会会長